

Encounters between the Elderly and Law Enforcement:  
An Overview of Mental Illness, Addictions, Victims, and Criminals

Mkay Bonner

University of Louisiana at Monroe

Mark S. Johnson

University of Louisiana at Monroe

Authors Note

Mkay Bonner, Departments of Criminal Justice and Psychology, University of Louisiana at Monroe.

Mark S. Johnson, University Police Department and Department of Criminal Justice, University of Louisiana at Monroe.

Correspondence concerning this article should be addressed to Mkay Bonner, Departments of Criminal Justice & Psychology, University of Louisiana at Monroe, Monroe, LA 71209. Contact: [bonner@ulm.edu](mailto:bonner@ulm.edu)

### Abstract

The elderly population is growing dramatically throughout the world. Out of this growth comes an increase in the number of encounters between the aged and law enforcement. These encounters occur because of a variety of factors including mental illnesses and addictions. Furthermore, older adults may be victims as evidenced in different forms of abuse such as physical, emotional, or financial abuse. Sadly, some documentation exists that older adults are committing more crimes. Multiple reasons have been postulated for these crimes including poverty, jealousy, and boredom. All of these different situations with the aged have created an increase in the number of encounters between the elderly and law enforcement. Different programs exist that can be used to train law enforcement professionals to identify and help the elderly who are in need of assistance. Regardless of the program used, law enforcement can improve their ability to protect and serve this vulnerable and growing segment of the population. The elderly need the help of law enforcement and this need will continue to increase as the elderly population grows.

*Keywords:* elderly, aging, geriatric, law enforcement, police, victims, criminals, mental illness, addictions, community programs, crisis intervention team training

### Encounters between the Elderly and Law Enforcement:

#### An Overview of Mental Illness, Addictions, Victims, and Criminals

Obvious to many, the aging population is increasing dramatically in numbers and percentages across the world (Kettaneh, 2015; Rutherford, 2012; Shrestha & Heisler, 2011; TRIP, 2018; United Nations, 2012). Professionals estimate that people over 65 years of age will constitute 20% of the population by 2020 and will continue to increase after that (Bourns, 2000; Lavretsky, 2018). With the increases in the aged population, it is prudent to expect an increase in the encounters between the elderly and law enforcement. These encounters will occur in many manifestations. The specific encounters briefly addressed herein include those related to mental illness, addictions, victims, and criminals. To complicate the situation, these manifestations may also have considerable overlap such as the mentally ill person who is the victim of a crime or the opioid-addicted individual who steals to survive. It is important to note that this is a compendium of issues and it is not an exhaustive presentation of all of the possible encounters between the elderly population and law enforcement professionals.

The parameters of the aging population differ from study to study and organization to organization (Acierno et al., 2010; TRIP, 2018; World Health Organization, 2018a). Typically, 65 and older is considered the standard. With the increase in the human lifespan, some researchers specified 75 or 80-years-old as the point of significant aging. The literature related to older persons in jail often considered 50-years-old as the beginning point of the aging population because of the severity and difficulties of life behind bars (please see the article by Hanser and Hu in this special issue on Aging in the *Contemporary Southern Psychologist* for a more detailed discussion of the elderly who are incarcerated). The literature utilized herein primarily documented 65 and older as the age group of interest. A few articles that are

mentioned, incorporated 60 to 64-year-olds as well and these articles were included because of the depth of helpful information they provided.

### **Mental Illness**

According to the World Health Organization (2018b), there is a worldwide increase in the number of persons suffering from a mental illness. These individuals are often referred to as consumers (Bonner & Johnson, 2013). The estimates suggested that 15% of elderly individuals suffer from a mental illness (World Health Organization, 2017). Despite the increase in consumers, there has been a drastic decrease in affordable services and available mental health beds at hospitals (Bonner & Johnson, 2013; Louisiana POST Council, 2018; Torrey, 1997). As a result, this trifecta of situations provides a forewarning of increased encounters between law enforcement officers and the elderly whether they are consumers, victims, addicts, criminals, or a combination of these (Bonner & Johnson, 2013; Cordner, 2006; Louisiana POST Council, 2018).

The most common mental disorders in the elderly are depression at 7% and dementia at 5% (World Health Organization, 2017). Girgus, Yang, and Ferri (2017) found that gender differences in depression rates continue into old age with women diagnosed with depression more than men are. Depression is not a requirement of getting older and should never be considered normal. Unfortunately, depression in the elderly is often ignored by primary care treating professionals. To improve the quality of life of elderly patients, treating professionals should always assess for depression. A proactive program including social supports should be utilized to reduce the occurrence and debilitating effects of depression in the elderly. Law enforcement encounters with elderly individuals who are depressed may be tragic because it may be in the form of attempted or completed suicide. Some of the depressed elderly individuals may also resort to suicide by cop.

Dementia occurs in more than one form but the most recognizable may be Alzheimer's dementia. The World Health Organization (2017) estimated that more than 50 million people suffer from some form of dementia and Alzheimer's dementia accounts for approximately 70% of these cases. Alzheimer's dementia involves cognitive impairment, which is manifested in problems with memory, thinking, and acting (Alzheimer's Association, 2018). Treatments include medications, environmental manipulations, caregiver training, and support groups. It is common for individuals with Alzheimer's dementia to wander and become lost. At this point, family members may contact the police to help find their loved one or the police may find the person wandering and try to assist in getting them home. In these cases, the goal of law enforcement involvement is to help the individual get home before a tragedy occurs.

In the aged, anxiety disorders occur nearly as often as dementia at a rate of almost 4% (World Health Organization, 2017). However, anxiety disorders do not get the attention that Alzheimer's dementia garners. Not surprisingly, anxiety was found to impact memory in older adults (Herrera, Montorio, & Cabrera, 2017). These authors also found older adults with anxiety remembered more events with negative associations than with positive associations. As with other disorders, anxiety in the elderly should be actively screened and treated. Disorders and disabilities should not be accepted without treatment just because someone is over 60-years-old. Typically, law enforcement officers are not called for help for someone with an anxiety disorder only. However, when an anxiety disorder is present with another disorder such as depression, the possibility of an encounter with law enforcement increases.

Psychotic disorders including schizophrenia and delusional disorders involve distorted thinking and loss of contact with reality. Most schizophrenic patients are younger in part because individuals with this disorder do not tend to have a long life. In elderly patients,

diagnosing and treating these disorders can be particularly problematic because of interrelated and compounding issues of a neurological, biological, and psychosocial nature (Mendez, von Gunten, & Antunes, 2015). One researcher succinctly stated, “Still too often, older patients with schizophrenia are seen as lost cases, although they should deserve our respect for having survived into old age with an illness as severe and disruptive as schizophrenia.” (Meesters, 2017, pp. 511). Crisis Intervention Team (CIT) police officers routinely answer calls for service that involve someone with schizophrenia. In keeping with the low prevalence rates, only approximately 5 – 10% of these calls involve an elderly individual (M. Fendall, personal communication, October 18, 2018). However, these calls still have the potential for danger for the elderly individual and the police.

Early diagnosis is important with any disorder including mental disorders (Alzheimer’s Association, 2017; Mendez, von Gunten, & Antunes, 2015; World Health Organization, 2017). Education about the disorder is critical for the individual with the diagnosis, family members, and/or other caregivers. The World Health Organization identified several disorders as needing improvements in programming including depression, psychoses, dementia, and substance use disorders. Improvements in these areas can produce benefits for the elderly and may help reduce their negative encounters with law enforcement.

### **Addictions**

The World Health Organization (2017) cited 1% of the worldwide elderly population suffer from some form of substance abuse. Importantly, they also stated that the misuse of substances in the elderly is often ignored or incorrectly diagnosed. This cited percentage appears relatively miniscule. However, when considering the size of the elderly population, this percentage equates to millions and millions of individuals who need help.

The National Survey on Drug Use and Health provides a plethora of information including some age-related characteristics (SAMHSA, 2017). The data are divided into lifetime use, past year use, and past month use. In addition, the detailed age characteristics include ages 60 to 64 and 65 and above. Many researchers utilize this data as they investigate specific areas of drug use. The most recent survey provided information through the year 2017.

According to this national survey (SAMHSA, 2017); the most commonly used drug by individuals 60 and older was alcohol. Over 60% of those surveyed admitted consuming alcohol during the past year. More importantly, almost 20% of 60 to 64-year-olds and over 11% of individuals aged 65 and older admitted to binge drinking during the past month. Furthermore, almost 5% of 60 to 64-year-olds and almost 3% of individuals 65 and older admitted to heavy drinking in the past month. Binge drinking was defined as five drinks for males and four drinks for females. Heavy drinking was defined as binge drinking more than five days in the past month. All of the percentages were higher for the 60 to 64 age group than the 65 and older age group.

Alcohol consumption by itself is not illegal for these age groups. It is acceptable in most jurisdictions in the U.S. once a person has reached the legal age of consumption, which is usually age 21. Often when people think about alcohol abuse, they think of young adults or possibly middle-aged men. In 2015, Kettaneh warned that alcohol abuse is not a disorder reserved for the young. Older adults suffer from alcoholism as well. They also have unique problems and complications as a result of the aging process combined with alcoholism including hypertension, falls, and dementia. Kettaneh made a strong case for counselors to become educated and proficient in treating alcohol abuse within the elderly population.

Reinforcing the concern of elderly alcoholism, Dombrowski, Norrel, and Holroyd (2016) documented alcohol abuse as the primary substance misused by elderly psychiatric patients. During a 10-year-period, they found approximately 12% of elderly psychiatric patients abused a substance with almost 75% of them abusing alcohol. From this elderly patient total, they found 11% addicted to sedative hypnotics, almost 3% addicted to opioids, and approximately 1% addicted to cannabis or tobacco. Almost 40% involved an unspecified addiction.

The progressive legalization of marijuana, (i.e. cannabis), may have had an impact on older adults as well as younger adults. In 2017, an article was published which analyzed the demographic trends of cannabis users from 2006 through 2013 (Han et al., 2017). The authors found a significant increase in cannabis usage with persons aged 65 and older. From 2006 to 2013, the cannabis usage increased 250%. This significant increase should forewarn professionals to become knowledgeable and competent to treat cannabis addictions in the elderly. This increase is also particularly problematic for law enforcement because of the contradictions in laws between some states and the federal government.

A recent study documented the types of substances that were consumed by the elderly who had been referred for addiction treatment (Saber, FaniSaben, Mousavinasab, Zarghami, & Taraghi, 2018). All of the participants who had been referred for addiction treatment had abused opioids. Of these people, 65% admitted to using tobacco products and 31% drank alcohol. Almost 27% of them admitted to using some form of sedative.

The opioid epidemic is abundant within younger age groups but the elderly are not immune. Psychiatrists Suryadevara, Holbert, and Averbuch (2018) bemoaned the drastic increase in opioid prescriptions, abuse, and overdoses within the aging population. However, they did not recommend for physicians to quit prescribing opioids to elderly patients entirely.



Instead, they stated the following: “We advocate strongly for vigilance in oversight, monitoring closely for warning signs of misuse, and thorough documentation.” (pp. 16). They emphasized that 80% of elderly long-term care patients have significant pain. They clearly stated there are appropriate situations to prescribe opioids for pain management, especially in end of life (hospice) care.

Research and review articles have not focused on the use of tobacco products recently as much as they did in the past. However, SAMHSA continues to include the use of tobacco products in the National Survey (SAMHSA, 2017). Tobacco products include cigarettes, cigars, and snuff. Between 66 to 71% of elderly respondents admitted to tobacco use at some point in their lives. Of particular interest is the past year and past month usage of tobacco products in the elderly. The survey documented a past year usage of 14 – 22% and a past month usage of 12 – 20%. As with other substances, tobacco can complicate health conditions especially in the elderly. Tobacco cessation should be a consideration even in the geriatric population. The elderly are unlikely to encounter the police because of the use of tobacco. However, it is informative that some elderly individuals continue to use tobacco products. Often CIT officers will utilize cigarettes as a bargaining tool. With the confirmation that some older adults continue to use tobacco, CIT officers can continue to use these products as they attempt to deescalate crisis situations.

Addictions in the elderly are not limited to drug substances only. Gambling is another addiction of concern (eSauvaget et al., 2015). The prevalence estimates of a gambling disorder in the elderly ranges to a high of almost 11%. The elderly primarily choose casinos and bingo for their gambling activities. As of 2015, online gambling had not been identified or researched within the elderly population. Nevertheless, eSauvaget and associates (2015) did identify an 83-

year-old individual with an online gambling disorder who began his online gambling habits when he was 80-years-old. The authors proposed that this is not an anomaly. They postulated that online gambling is an under-reported and under-investigated phenomenon. The common physical problems inherent with aging such as mobility difficulties make online gambling an easy option for older adults. Many people over 60 may choose to gamble online because of these physical problems, loneliness, and isolation. Sadly, engaging in online gambling will only exacerbate problems of isolation and depression. The authors encouraged research and development of age-specific treatments for online gambling addictions in the elderly.

Interestingly, circadian rhythms may provide a key to addiction treatments in the young and the aged (Gulick & Gamsby, 2018). The human bio-clock has provided another avenue for addiction treatment as it now appears to influence the reward systems of drugs. Research is proving beneficial in this area.

However, the largest gap in knowledge is in our understanding of how age mediates the interactions of the circadian and reward systems. Melatonin levels, chronotype, and addiction patterns all vary with age, but almost all work has been done in young adults, and in adult animal model systems. More recently, research has begun to examine the importance of the adolescent night owl chronotype in addiction, but more studies are needed, and almost no work has been done in elderly adults or comparable model systems (Gulick & Gamsby, 2018, pp. 137).

This information reinforces the need for more research and evidence-based treatment for addictions within the elderly population.

In the sample of 17,000 elderly persons identified by Ghossoub and Khoury (2018), slightly over 2% had a substance use disorder as compared to the approximately 12% who had a

psychiatric disorder. This percentage regarding substance abuse is double the World Health Organization's (2017) estimations. Within this 17,000-person sample, almost 11% had received mental health treatment in the past year but only 0.1% had received alcohol abuse treatment. The previously mentioned National Survey contained similar results but they identified that approximately 2% of elderly individuals had received some form of substance abuse treatment during the past year (SAMHSA, 2017). This lack of substance abuse treatment for the elderly is troubling. This overall situation of addictions makes some elderly individuals more vulnerable to negative encounters with law enforcement.

### **Victims**

The elderly can be particularly attractive targets for criminals. They are a vulnerable population who may not have the knowledge, cognitive abilities, or physical strength to avoid becoming a victim of crime. In a cursory review of FBI databases, the most frequent articles associated with law enforcement and the elderly focused on fraud. Many types of fraud exist including home improvement frauds and identity theft. Acierno and colleagues (2010) found over 5% of elderly respondents admitted being the victim of financial fraud committed by family members. According to the 2017 revised bulletin on identify theft, the number of elderly victims increased from 5% to almost 6% between 2012 and 2014 (Harrell, 2017). They were the only age group to have a significant increase during this timeframe. The vast majority of these cases involved the misuse of existing credit cards and bank accounts. Unfortunately, most of these cases are never reported to the police.

Elder abuse is a crime. The American Association of Retired Persons (AARP) has warned about elder abuse estimating 10% of older adults have been victimized (Schmitt, 2017). They especially warned about abuse from hired caregivers. In 2010, the National Elder

Mistreatment Study was conducted in the U.S. (Acierno et al., 2010). The authors emphasized the prevalence of many forms of elder abuse including physical, financial, and sexual abuse. Their results documented the following: almost 5% experienced emotional abuse, slightly under 2% reported physical abuse, and less than 1% claimed they were sexually abused. Greater than all of these was the percentage of elderly individuals who reported being neglected during the previous year, which was over 5%.

The National Center on Elder Abuse (NCEA, 2018) is a government-supported resource through the U.S. Administration on Aging. The NCEA (2018) also cited that 10% of senior citizens were the victim of elder abuse. Based on census data, this percentage equates to over 4 million elderly adults who are victims. Even more disturbing, Acierno and associates (2010) found that 50% of elderly persons with dementia would be the victim of some form of abuse.

The NCEA (2018) was clear in stating that elder abuse is underreported. The World Health Organization (2017) who estimated that almost 16% of elderly individuals suffer some form of elder abuse reinforces this previously. They also believed that this percentage is an underestimation. The World Health Organization (2018a) stated that elder abuse occurs more in residential facilities than in other environments. In addition, they do anticipate increases in elder abuse throughout the world as the elder population increases.

The website for the NCEA (2018) provides many resources and fact sheets to help prevent elder abuse or to assist if abuse has occurred. Their resources address a variety of areas including crime, neglect, disabilities, and domestic violence. The World Health Organization (2018a) also promotes training for providers of elderly services, respite for family caregivers, and public and professional awareness campaigns. Adult Protective Services are available in

most states. Moreover, law enforcement agencies usually have divisions or officers specifically designated to work with elderly citizens.

The elderly may also be victims in motor vehicle accidents, even if they are the ones driving. The number of people who are over 65 and driving in the U.S. increased by almost 40% from 2006 to 2016 (TRIP, 2018). The largest increases in drivers over 65 were found in Louisiana at 44%, Arkansas at 28% and South Carolina at 28%. According to the most recent statistics, there have been increases of elderly drivers of 20% or more in 19 states. With more drivers, there are more accidents. The elderly are especially vulnerable in car crashes due to poorer overall health and fragility. The geriatric driving population has the tendency to travel at off-peak times. However, with increased traffic congestion, the times that are available for them to drive become less and less. Unfortunately, elderly men and women tend to live beyond their driving abilities by 7 to 10 years. Transportation mobility makes a tremendous impact on the elderly population's quality of life. Once they cannot drive, their interactions with others decreases greatly. According to TRIP, there are ways to improve safety for geriatric drivers. Vehicle technology such as blind spot warnings helps the elderly avoid accidents. Roads that are in good repair with adequate markings and sufficient lighting are important. In addition, some states have found better results when elderly drivers are required to renew their driver's licenses in-person. Many of these individuals no longer seek a license and stop driving of their own volition. Law enforcement may help the families of older adults by referring the elderly individual to the Department of Motor Vehicles if driving ability is of concern.

### **Criminals**

While the elderly are more often viewed as victims, they may also commit criminal actions. Speiser (2015) wrote a business article entitled "The World's Elderly are on a Crime

Spree”. One specific major heist of jewelry and money involved nine thieves ages 43 through 76. Speiser promulgated that the aged respected the law in the past but now they demonstrate a growing disregard. He stated that the commission of crimes is on the increase among older individuals in the U.S. and throughout the world. South Korea experienced a 40% increase in violent crimes committed by senior citizens between 2011 and 2013. In Japan, crimes by the elderly have doubled since 2003. In England, the incarceration rates of the geriatric population increased three times more than the rates for other age groups in the general prison population. Speiser attributed these increases to inadequate income, increased expenses especially in food and medical costs, boredom, and jealousy of other affluent elderly persons.

One recent article entitled “Granny Dealing Drugs on the Government’s Dime: Why Medicare and Medicaid Should Have Safeguards in Place to Prevent Abuse” addressed the elderly as criminals (Campbell, 2017). In this article, Campbell emphasized the increased ease with which elderly patients can receive prescriptions for pain medications. Often they can receive large quantities of these medications, which is not possible for younger patients. In addition, Medicare and/or Medicaid do not closely monitor these prescriptions. Compounding the problem is the fact that most of these elderly patients live on a small, fixed income. These factors set the stage for elderly drug dealers. Extrapolating estimates from the U.S. Government Accountability office, approximately 650,000 elderly individuals across the U.S. may be obtaining prescription medications to sell for profit. Interesting, Louisiana is one of the few states with mandatory disclosure laws, which attempt to prevent patients from doctor shopping for prescriptions. In conclusion, Campbell made a plea for better oversight and management of these medications with the elderly to reduce fraud and protect taxpayers’ money.

In further consideration of the elderly as criminals, Ghossoub and Khoury (2018) conducted several analyses based on the National Survey on Drug Use and Health from 2006 through 2014. "...there is increasing evidence that older adults are committing more offenses and violent crimes leading to an increased frequency of arrests in late life" (pp. 211). The sample size of people who were 65 and older was impressive at more than 17,000 individuals. Of these, only 5% admitted to committing a criminal offense or being arrested during the previous year. Drunk driving was found to be the most common criminal offense by the elderly with a prevalence rate of 18% of the total reported arrests. Similarly, 12% were arrested for being drunk. Additional arrests of interest included vehicle theft at 8%, burglary at over 6%, and larceny and other assaults both reached 5.8%. Serious violent offenses reached 5.6% and was defined as "aggravated assault, rape, homicide, or manslaughter" (Ghossoub & Khoury, 2018, pp. 214). Previously, Campbell (2017) stressed the concern of the aged defrauding the government by obtaining prescription medications to be sold for profit. Ghossoub and Khoury (2018) found that 3.9% of the elderly population who admitted to being arrested during the past year were arrested for the possession and sales of drugs. Obviously, with more criminal activity, there will be more encounters between the police and the elderly.

### **The Law Enforcement Response**

In 2000, Bourns emphasized the importance of police providing services for the elderly. He stressed the growing elderly population and the value of a different approach for law enforcement and elderly interactions. Law enforcement officers should be trained in methods to address elder abuse and protocols for Alzheimer's patients. In 2014, this call is reiterated (Brown, Ahalt, Steinman, Kruger, & Williams, 2014). These authors emphasized the police as the front line for geriatric healthcare in the community. Included in their study were all calls for

service with the elderly including first responder calls related to health conditions. In this research, over 80% of the law enforcement officers stated they had been trained in issues relating to the elderly. However, only 32% considered themselves knowledgeable in this area. The police requested more training in elderly health conditions and in available community resources.

In a follow-up study, Brown et al. (2017) evaluated a training program for police that focused on geriatric issues. This short training program was associated with a comprehensive CIT program. Forty-five percent of officers who attended the training reported having daily interactions with the elderly. According to post-test scores, knowledge of aging issues did increase. Through this brief two-hour training, officers felt more prepared to respond to elderly individuals.

Since 2006, law enforcement officers throughout Louisiana have been taught about mental illnesses including some elderly conditions (Bonner & Johnson, 2013; Region VIII Mental Health Community Response Task Force, 2010). In northern Louisiana, the primary program has been conducted as part of the Northeast Delta Crisis Intervention Team Training Programs. This 40-hour training program provides law enforcement officers with knowledge and experiential training on mental illnesses, substance abuse disorders, and Alzheimer's dementia. Over the 12 years since its inception, this program has conducted almost 50 of the weeklong trainings and trained over 1000 officers, dispatchers, and other public safety personnel (Bonner & Johnson, 2018). This CIT program has also sponsored continuing education in elder issues such as the Silver Alert program. The course evaluations, post-test scores, and anecdotal stories over the years have substantiated the value of this program for training officers in a myriad of crisis situations including encounters with the elderly.



Hamm, Wylie, and Brank (2018) researched the level of confidence aging persons had in law enforcement and the courts. They linked positive perceptions with cooperation and compliance, both of which are important to successful encounters with the police. According to their research, they found that older individuals documented a greater trust in law enforcement than younger individuals. They suggested that further research is warranted before statements can be made to suggest that this trust and confidence will increase crime reporting by the elderly.

### **Conclusions**

The evidence is substantial: The elderly population is growing. It is not unreasonable to expect an increased number of encounters with law enforcement as these geriatric numbers increase. Some sources emphasized deinstitutionalization as the main reason there are more encounters between the police and citizens of any age (Cordner, 2006; Louisiana POST Council, 2018; Torrey, 1997). This may be accurate for individuals with a mental illness or addiction but it does not explain all of the possible reasons for an increase in encounters with law enforcement. It is not uncommon for the elderly to be a victim of a crime and need police assistance. And, some articles have documented the increase in the number of elderly persons as criminals. All of these situations elucidate the main point: The elderly population is growing and will continue to do so. Finding ways to help the geriatric population get the help they need will be one important step toward reducing negative encounters with the police.

A recurrent theme with the elderly is that conditions and disorders are often ignored, misdiagnosed, or inappropriately treated. Education is critical to combat these failures. Elderly persons and their caregivers need to be taught regarding what is normal and abnormal in the aging process. Treating professionals need good training related to the elderly and the manifestation of disorders specific to the aging population. Public awareness campaigns will

also be helpful. Law enforcement professionals need training about the elderly including information on identifying behaviors related to illnesses and addictions, victim characteristics, and criminal manifestations. More evidence-based research is needed to guide all of this education.

With the increase in numbers of the elderly population and the critical absence of appropriate resources, there will be no lack of customers, clients, or patients for the treating professional who chooses to specialize in this demographic. If someone wants their work to make a difference, a career serving the elderly population should prove to be both satisfying and rewarding.

The ultimate goal for law enforcement professionals is to protect and serve the citizens. The elderly are one of the most vulnerable subgroups of these citizens. Additional knowledge regarding disorders and victimology among the elderly will enhance the service that law enforcement provides to the aged. Increasing the positive encounters between law enforcement and the elderly is an admirable and respectable goal.

## References

- Acierno, R., Hernandez, M., Amstadter, A., Resnick, H., Steve, K., Muzzy, W., & Kilpatrick, D. (2010). Prevalence and correlates of emotional, physical, sexual, and financial abuse and potential neglect in the United States: The national elder mistreatment study. *American Journal of Public Health, 100*(2), 292-297.
- Alzheimer's Association. (2018). 2018 Alzheimer's Disease facts and figures. *Alzheimer's Dementia, 14*(3), 367-429.
- Bonner, M., & Johnson, M. (2013). The intersection between law enforcement and persons with a mental illness. *Crime, Punishment, and the Law, 3*(1), pp. 15-26.
- Bonner, M., & Johnson, M. (2018). The examination of the techniques utilized during crisis incidents based on CIT officer incident reports. Paper presented at the 44<sup>th</sup> Annual Conference of the Society for Police and Criminal Psychology, Sarasota, FL.
- Bourns, W. (2000). Police gerontology services for the elderly: A policy guide. *Justice Professional, 13*(2), 179-193.
- Brown, R., Ahalt, C., Rivera, J., Stijacic Cenzer, I., Wilhelm, A., & Williams, B. (2017). Good cop, better cop: Evaluation of a geriatrics training program for police. *Journal of the American Geriatrics Society, 65*(8), 1842-1847.
- Brown, R., Ahalt, C., Steinman, M., Kruger, K., & Williams, B. (2014). Police on the front line of community geriatric health care: Challenges and opportunities. *Journal of the American Geriatrics Society, 62*(11), 2191-2198.
- Campbell, K. (2017). Granny dealing drugs on the government's dime: Why Medicare and Medicaid should have safeguards in place to prevent abuse. *Elder Law Journal, 24*(2), 401-428.

- Cordner, G. (2006). *People with a mental illness*. Community Oriented Policing Services; Problem-Oriented Guides for Police. The U.S. Department of Justice.
- Dombrowski, D., Norrell, N., & Holroyd, S. (2016). Substance use disorders in elderly admissions to an academic psychiatric inpatient service over a 10-year period. *Journal of Addiction, 2016*. doi: 10.1155/2016/4973018
- eSauvaget, A., eJimenez-Murcia, S., eFernandez-Aranda, F., eFagundo, A., eMoragas, L, eWolz, I., ... Menchon, J. (2015). Unexpected online gambling disorder in late-life: A case report. *Frontiers in Psychology, 6*, doi: 10.3389/fpsyg.2015.00655
- Ghossoub, E., & Khoury, R. (2018). Prevalence and correlates of criminal behavior among the non-institutionalized elderly: Results from the national survey on drug use and health. *Journal of Geriatric Psychiatry, 31*(4), 211-222.
- Girgus, J., Yang, K., & Ferri, C. (2017). The gender difference in depression: Are elderly women at greater risk for depression than elderly men? *Geriatrics, 2*(4), 35.
- Gulick, D., & Gamsby, J. (2018). Racing the clock: The role of circadian rhythmicity in addiction across the lifespan. *Pharmacology & Therapeutics, 188*, 124-139.
- Hamm, J., Wylie, L., & Brank, E. (2018). Measuring older adult confidence in the courts and law enforcement. *Criminal Justice Policy Review, 29*(3), 301-316.
- Han, B., Sherman, S., Mauro, P., Martins, S., Rotenberg, J., & Palamar, J. (2017). Demographic trends among older cannabis users in the United States, 2006-13. *Addiction, 112*(3), 516-525.
- Harrell, E. (2017). *Victims of Identity Theft, 2014*. Bureau of Justice Statistics Bulletin. U.S. Department of Justice.

- Herrera, S., Montorio, I., & Cabrera, I. (2017). Effect of anxiety on memory for emotional information in older adults. *Aging & Mental Health, 21*(4), 362-368.
- Kettaneh, A. (2015). Substance abuse among the elderly population: Overview and management. *Journal of Applied Rehabilitation Counseling, 46*(4), 11-17.
- Lavretsky, H. (2018). Geriatric psychiatry introduction: Neuropsychiatric disorders of aging. *Psychiatric Times, 35*(1), 15-21.
- Louisiana POST Council. (2018). *The Louisiana POST Police Officer Training Manual*. Baton Rouge, Louisiana: Louisiana POST Council.
- Mendez, M., von Gunten, A., & Antunes, M. (2015). Schizophrenias and delusional disorders amongst the elderly. *Swiss Archives of Neurology, Psychiatry and Psychotherapy, 166*(4), 113-125.
- Meesters, P. (2017). Healthy older schizophrenia patients: Exceptions to the rule? *The American Journal of Geriatric Psychiatry, 25*(5), 510-511.
- National Center on Elder Abuse. (2018). U.S. Administration of Aging. Retrieved from <https://ncea.acl.gov/>
- Region VIII Mental Health Community Response Task Force. (2010). *Northeast Delta CIT Training Manual*. Monroe, LA: Region VIII Mental Health Community Response Task Force.
- Rutherford, T. (2012). Population ageing: Statistics (Report to the House of Commons, pp. 1-8). Retrieved from [www.parliament.uk/briefing-papers/sn03228.pdf](http://www.parliament.uk/briefing-papers/sn03228.pdf)
- Saberi, M., FaniSaben, L., Mousavinasab, N., Zarghami, M., & Taraghi, Z. (2018). Substance use disorders in the elderly people referring to addiction treatment clinics, 2017. *Directory of Open Access Journals, 6*(2), 31-40.

SAMHSA. (2017). *National survey on drug use and health, 2017*. Center for Behavioral Health Statistics and Quality. Substance Abuse and Mental Health Services Administration (SAMHSA), U.S. Department of Health and Human Services.

Schmitt, R. (2017). Elder abuse: When caregiving goes wrong. *AARP Bulletin*. American Association of Retired Persons. Retrieved from <https://www.aarp.org/caregiving/basics/info-2017/elder-abuse-assisted-living.html>

Shrestha, L. B., & Heisler, E. J. (2011). The changing demographic profile of the United States (Report for Congress, pp. 1-36). Congressional Research Service. Retrieved from <http://www.fas.org/sgp/crs/misc/RL32701.pdf>

Speiser, M. (2015, May 30). The world's elderly are on a crime spree. *Business Insider*. Retrieved from [www.businessinsider.com/the-elderly-are-on-a-crime-spre-2015-5](http://www.businessinsider.com/the-elderly-are-on-a-crime-spre-2015-5)

Suryadevara, U., Holbert, R., & Averbuch, R. (2018). Opioid use in the elderly. *Psychiatric Times*, 31(1), 15-21.

Torrey, E. (1997). *Out of the shadows*. NY, NY: Wiley.

TRIP. (2018). *Preserving the mobility and safety of older Americans*. National Transportation Research Group. Washington, D.C.

United Nations. (2012). *Population aging and development*. Available from [www.unpopulation.org](http://www.unpopulation.org)

World Health Organization. (2017). *Mental health of older adults*. Fact Sheet. The World Health Organization. Retrieved from <http://www.who.int/en/news-room/fact-sheets/detail/mental-health-of-older-adults>

World Health Organization. (2018a). *Elder Abuse*. Fact Sheet. The World Health Organization. Retrieved from <http://www.who.int/en/news-room/fact-sheets/detail/mental-disorders>

World Health Organization. (2018b). *Mental Health*. Fact Sheet. The World Health Organization. Retrieved from <http://www.who.int/en/news-room/fact-sheets/detail/mental-disorders>